

Up-to-date Questions and Answers from authentic resources to improve knowledge and pass the exam at very first attempt. ----- Guaranteed.



Podiatry-License-Exam-Part-III Dumps Podiatry-License-Exam-Part-III Braindumps Podiatry-License-Exam-Part-III Real Questions Podiatry-License-Exam-Part-III Practice Test Podiatry-License-Exam-Part-III Actual Questions



killexams.com

APMLE

Podiatry-License-Exam-Part-III

NBPME Podiatry License Exam Part III - 2025







What is the most common cause of a chronic turf toe?

- A. Hyperextension of the hallux
- B. Trauma
- C. Infection
- D. Neurological disorder

Answer: A

Question: 310

What is the most appropriate treatment for a patient with a chronic proximal phalanx dislocation?

- A. Physical therapy
- B. Surgery
- C. Immobilization
- D. Rest and ice

Answer: B

Question: 311

What is the most effective treatment for a patient with a chronic Morton's neuroma?

- A. Physical therapy
- B. Steroid injection
- C. Surgery
- D. Rest and ice

Answer: C

Question: 312

What is the most common cause of a chronic plantar plate tear?

- A. Overuse
- B. Trauma
- C. Infection
- D. Neurological disorder

Answer: A

Question: 313

What is the most appropriate treatment for a patient with a chronic fifth metatarsal stress fracture?

- A. Immobilization
- B. Physical therapy
- C. Steroid injection
- D. Surgery

Answer: A



Question: 314

What is the most effective treatment for a patient with a chronic medial tibial stress syndrome?

- A. Physical therapy
- B. Immobilization
- C. Surgery
- D. Rest and ice

Answer: B

What is the most common cause of a chronic interdigital neuroma?

- A. Overuse
- B. Trauma
- C. Infection
- D. Neurological disorder

Answer: A

Question: 316

What is the most appropriate treatment for a patient with a chronic talus fracture?

- A. Physical therapy
- B. Surgery
- C. Immobilization
- D. Rest and ice

Answer: B

Question: 317

What is the most effective treatment for a patient with a chronic sesamoid stress fracture?

- A. Physical therapy
- B. Immobilization
- C. Surgery
- D. Rest and ice

Answer: B

What is the most common cause of a chronic lateral column overload syndrome?

- A. Overuse
- B. Trauma
- C. Infection
- D. Neurological disorder

Answer: A

Question: 319

What is the most appropriate treatment for a patient with a chronic Lisfranc dislocation?

- A. Physical therapy
- B. Surgery
- C. Immobilization
- D. Rest and ice

Answer: B

Question: 320

What is the most effective treatment for a patient with a chronic hallux limitus?

- A. Physical therapy
- B. Steroid injection
- C. Surgery
- D. Rest and ice

Answer: C

What is the most common cause of a chronic medial column overload syndrome?

- A. Overuse
- B. Trauma
- C. Infection
- D. Neurological disorder

Answer: A







For More exams visit https://killexams.com/vendors-exam-list

